

Vaccine Information

The following vaccines require multiple doses over time. Provide the date your child received each dose.

Vaccine	Vaccine Schedule	Date Child Received Vaccine
Hepatitis B	Birth (first dose)	
	1-2 months (second dose)	
	6-18 months (third dose)	
Rotavirus	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
Diphtheria, Tetanus, Pertussis	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
	15-18 months (fourth dose)	
	4-6 yearsn (fifth dose)	
Haemophilus Influenza Type B	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
	12-15 months (fourth dose)	
Pneumococcal	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
	12-15 months (fourth dose)	
Inactivated Poliovirus	2 months (first dose)	
	4 months (second dose)	
	6-18 months (third dose)	
	4-6 years (fourth dose)	
Influenza	Yearly, starting at 6 months. Two doses given at least four weeks apart are recommended for children who are getting the vaccine for the first time and for some other children in this age group.	
Measles, Mumps, Rubella	12-15 months (first dose)	
	4-6 years (second dose)	
Varicella	12-15 months (first dose)	
	4-6 years (second dose)	
Hepatitis A	12-23 months (first dose)	
	The second does should be give six-8 months after first dose.	

For additional information about immunizations, visit the Texas Department of State Health Services website at www.dshs.state.tx.us/immunize/public.shtml

HHSC values your privacy. For more information, read our privacy policy online at <https://hhs.texas.gov/policies-practices-privacy#security>

Physcial or Public Health Personnel Verification

Signature or stamp of a physician or public health personnel verifying immunization information above:

Signature

Date